

# Spruce and Peatland Responses Under Changing Environments

mnspruce.ornl.gov



An experiment to assess the response of northern peatland ecosystems to increases in temperature and exposures to elevated atmospheric CO<sub>2</sub> concentrations

Peatlands cover only 3% of Earth's land surface but contain about 30% of the global soil carbon pool. Peat deposits originated from higher plants and moss-generated litter. Because of cold, oxygen-poor conditions, the carbon contained in northern peatlands has accumulated for thousands of years. Under current warming trends and consistent with climate projections, such accumulations of carbon may be vulnerable to further decomposition or mineralization. If global temperatures warm as projected at higher latitudes, these peatlands could release large amounts of greenhouse gases such as carbon dioxide (CO<sub>2</sub>) and methane (CH<sub>4</sub>) that could accelerate global warming. The ability to predict or simulate the fate of the stored carbon in response to environmental disruption remains hampered by a limited understanding of the controls of carbon turnover and the composition and functioning of peatland ecosystems.

To identify and quantify these critical environmental response mechanisms, the Terrestrial Ecosystem Science (TES) program within the Department of Energy's (DOE) Office of Biological and Environmental Research (BER) is supporting a whole-ecosystem experiment in an ombrotrophic bog (i.e., a raised bog that receives all water and nutrients from direct precipitation) located in the Marcell Experimental Forest of northern Minnesota. The Spruce and Peatland Responses Under Changing Environments (SPRUCE) experiment project, led by Oak Ridge National Laboratory (ORNL), enables the assessment of ecological responses across multiple spatial scales—including microbial communities, moss populations, various higher plant types, and some insect groups. The project is evaluating a wide range of increased temperatures and levels of elevated atmospheric CO<sub>2</sub> concentrations. Direct and indirect effects of the experimental perturbations are being tracked and analyzed over a decade. This



**Aerial View of SPRUCE Project Site.** SPRUCE research is being conducted on an 8.1 hectare peatland of the Marcell Experimental Forest in northern Minnesota. Ten open-topped aboveground enclosures are being used to simulate various levels of warming and CO<sub>2</sub> exposure. The remote landscape includes a mix of uplands, bogs, fens, lakes, and streams.

comprehensive suite of spruce-peatland process studies and observations is being linked to model development and application requirements for improving process representation, calibrating models, and evaluating model predictions for boreal ecosystems. SPRUCE is a cooperative joint venture by scientists from DOE national laboratories, the U.S. Department of Agriculture's (USDA) Forest Service, and universities.

## Warming and CO<sub>2</sub> Treatments

By 2100, future terrestrial environments are projected to be 4°C to 8°C warmer than today, depending on the latitude. Mean deep soil (>1 m) temperatures will also rise with a warming climate. A series of large, open-topped aboveground enclosures and a new method for warming soils from the surface down to approximately 2 m were installed in the peatland to simulate various levels of warming and CO<sub>2</sub> exposure at the whole-ecosystem scale. Soil and air temperatures within these enclosures cover multiple levels of warming from ambient to +9°C. Simultaneously, atmospheric CO<sub>2</sub> levels within half of the enclosures are elevated up to 800 to 900 parts per million to reflect current expectations for the levels that may be associated with end-of-century temperatures.

This new experimental system provides a platform for testing the mechanisms controlling the vulnerability of organisms, biogeochemical processes, and ecosystem functions to important environmental change variables (e.g., thresholds of organisms to decline or mortality, limitations to regeneration, biogeochemical limitations to productivity, and the cycling and release of CO<sub>2</sub> and CH<sub>4</sub> to the atmosphere). The ultimate goal is to determine the levels of warming at which

## Key SPRUCE Science Questions

The experiment's overarching science questions cover ecosystem responses ranging from the microbe to landscape scale. They include:

- Will deep belowground warming in the future release 10,000 years of accumulated carbon from peatlands that store one-third of Earth's terrestrial carbon? At what rate?
- Will these carbon releases be in the form of CO<sub>2</sub> or CH<sub>4</sub> with 30 times the warming potential?
- Are peatland ecosystems and organisms vulnerable to atmospheric and environmental change? What changes are likely?
- Will ecosystem services (e.g., regional water balance) be compromised or enhanced by atmospheric and climatic change?

Answers to these questions will provide insights not only for small-scale processes but also for landscape-relevant water, carbon, and energy fluxes for similar peatlands. Results will inform higher-order models of vegetation responses under various levels of climatic warming and associated end-of-century atmospheric change.




U.S. DEPARTMENT OF  
**ENERGY**

Office of  
Science

Office of Biological and Environmental Research

[science.energy.gov/ber/](http://science.energy.gov/ber/)





**Whole-Ecosystem Warming Method.** *Experimental enclosures were built to provide warming from the tree tops to the deep soil (–2 m to –3 m). The open-topped enclosures encompass 12-m diameter internal study areas, with 8-m tall side walls. A subsurface corral isolates the belowground peat environment for measures of local hydrologic conditions. Aboveground warming enclosures became fully operational in 2015. Inset: Closeup of enclosure.*

ecosystems will reach a critical change in temperature and CO<sub>2</sub> levels that would push them into a new state (i.e., altered community composition and capacity to store carbon). Deep peatland heating was initiated in June 2014 to independently test the impacts of warming on the bog’s ancient deep carbon stocks, ahead of the full whole-ecosystem warming that began in June 2015.

### Connecting Observations to Models for Improved Earth System Predictions

New modeling approaches are needed to incorporate the complex relationships among warming, drying, mineralization processes, and vegetation responses associated with environmental changes. The experimental data generated by SPRUCE are providing quantitative evidence of the effects of elevated temperatures and CO<sub>2</sub> on northern peatland ecosystems and the vast carbon stores associated with the hydrology and biogeochemistry of these globally widespread landscapes. These data have led to key improvements in how biogeochemistry models (and further community models) represent the temperature dependence of carbon losses as CO<sub>2</sub> and CH<sub>4</sub> and the extent to which they might be counterbalanced by enhanced net primary production that is driven by a longer growing season, nutrient enrichment, and elevated atmospheric CO<sub>2</sub> levels.

### Research Platform for the Scientific Community


The core suite of SPRUCE research is being pursued by scientists at ORNL and the USDA’s Forest Service. Collaborations with 30 univer-

sities have been established to leverage the project. Also welcomed are new initiatives consistent with the design, science mission, spatial constraints, and integrity of the experiment on the sensitive bog ecosystem. Collaboration opportunities include but are not limited to:

- Canopy albedo changes
- Remote sensing of canopy function
- Microscale LIDAR
- Bole respiration
- Trace gas emissions
- Herbivory
- Food web linkages
- Pests and pathogens (host defenses versus pathogen virulence)

For more information on how to become involved with the SPRUCE project, go to [mnspruce.ornl.gov/contact/](http://mnspruce.ornl.gov/contact/).

**Picea-Sphagnum Bog.** *Picea mariana, or black spruce, and Sphagnum, a genus of peat mosses, are the dominant overstory and groundcover, respectively, of the boreal peatland where SPRUCE is being conducted. The bog contains deep peat deposits perched several meters above the regional groundwater table.*



**Environmental Monitoring.** *Left: Periodic shrub, Sphagnum, and peat community-level measurements of CO<sub>2</sub> and CH<sub>4</sub> are assessed from 1.2-m diameter in situ collars left in the experimental plots for seasonal and treatment response evaluations. Right: SPRUCE experimental plot showing the environmental monitoring mast and visible infrastructure for belowground gas flux and pore water measurements prior to the addition of the aboveground enclosure.*

### Program Manager and Websites

**Daniel B. Stover**, [daniel.stover@science.doe.gov](mailto:daniel.stover@science.doe.gov), 301.903.0289

**SPRUCE** [mnspruce.ornl.gov](http://mnspruce.ornl.gov)

**Terrestrial Ecosystem Science**  
[tes.science.energy.gov](http://tes.science.energy.gov)

**Climate and Environmental Sciences Division**  
[science.energy.gov/ber/research/cesd/](http://science.energy.gov/ber/research/cesd/)

**DOE Office of Biological and Environmental Research**  
[science.energy.gov/ber/](http://science.energy.gov/ber/)

**DOE Office of Science**  
[science.energy.gov](http://science.energy.gov)

**U.S. Department of Energy**  
[energy.gov](http://energy.gov)

*Images were provided by Oak Ridge National Laboratory.*